

A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.

Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.

LA-UR-84-1640

CONF-8405180--1

NOTICE

PORTIONS OF THIS REPORT ARE ILLEGIBLE. It has been reproduced from the best available copy to permit the broadest possible availability.

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA-UR--84-1640

DE84 012453

TITLE THEORETICAL STUDY OF CEPHEID LIGHT CURVES

AUTHOR(S) C. G. Davis

SUBMITTED TO IAU Colloquium #82, Toronto, Canada, May 29-June 1, 1984.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

MASTER

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

THEORETICAL STUDY OF CEPHEID LIGHT CURVES

C.G. Davis

Los Alamos National Laboratory, Los Alamos, New Mexico

First line of INTRODUCTION after opening

Starting with the initial understanding that pulsation in variable stars is caused by the heat engine of Hydrogen and Helium ionization in their atmospheres (A.S. Eddington in Cox 1980) it was soon realized that non-linear effects were responsible for the detailed features on their light and velocity curves. With the advent of the computer we were able to solve the coupled set of hydrodynamics and radiation diffusion equations to model these non-linear features (Christy 1968, Cox et. al. 1966). Calculations including the effects of multi-frequency radiative transfer (Davis 1971) showed that grey diffusion was adequate for modeling Cepheids but not for the RR Lyrae or W. Virginis type variables. In 1977, in collaboration with J. Castor and D. Davidson (1977), we developed a non-Lagrangian method to resolve the region of the ionization front and remove the zoning effects previously found in theoretical light curves (Keller and Mutschlecner 1972). The new code (DYN) has been used in recent studies by Takeuti (1983), with Simon (1983), in an attempt to understand modal coupling, and with Moffett and Barnes (1981) in a detailed study of X Cygni. A minor success for DYN was obtained by comparing "phase lags", as determined from the first order modal expansions between maximum light and maximum outward velocity, with some low period Cepheid observations (Simon 1984). Another support for the "dips" as observed by Moffett and Barnes (1984) is the observation by Schmidt and Parsons (1982) of a "dip" in the IUE spectra of δ Cep near the phase of 0.85.

In this paper we want to describe some recent model results for long period (LP) Cepheids in an attempt to understand these "dips" and possibly get another handle on Cepheid masses. In section II we discuss these results and in section III we consider the implications of these model results on the problem of the Cepheid mass discrepancy.

LONG PERIOD CEPHEID MODELS

It appears that a classification of Cepheids could be: low amplitude low period sinusoidal Cepheids, "bump" Cepheids with periods from 7 to 10 days, and long period Cepheids with periods longer than 10 days. In this paper we are considering this latter class of variables. The models we have selected have effective temperatures near the middle of the instability strip and nearly constant at 5300 or 5500 K. We are mainly interested in seeing the effects of mass on

the light curves so we selected near evolutionary masses and 60% or so of the evolutionary masses. The period range was picked in the range from 12 to 22 days with selected stars from the Moffett and Barnes list for comparisons (Table I).

ch

TABLE I

PERIOD (PROTOTYPE)	LONG PERIOD CEPHEID MODELS			HIGH MASS		
	LUM	TEFF	P ₂ /P ₀	LUM	TEFF	P ₂ /P ₀
12.9 (Z SCT)	1.5	4.5 M ₀ .54	.499	2.5	7.5 M ₀ .55	.516
14.8 (RW CAS)	2.1	5.0 M ₀ .551	.497	3.0	8.5 M ₀ .54	.527
16.4 (X CYG)	2.45	5.7 M ₀ .54	.496	3.34	9.0 M ₀ .53	.520
22.0 (WZ SGR)	3.5	6.0 M ₀ .54	.475	5.5	10.0 M ₀ .54	.507

LUM ($\times 10^{37}$ erg/cm²-s) TEFF ($\times 10^4$ °K)

First line of te

First we note, from Table I, that the low and high mass models straddle the linear theory resonance boundary parameter of $P_2/P_0=0.5$. Near 0.5 the resonance interaction is the strongest and the Christy "bump" should appear near maximum light. A reasonable explanation of the connection between resonance theory and the "echo" is given by Whitney (1983). In Fig. 1 we illustrate the DYN results of this phenomena for model of a 10 day Cepheid with evolutionary mass and 60% of evolutionary mass. In the low mass model the "echo", which causes the "bump" at the surface, can clearly be seen near the inner boundary.




Figure 1 - Velocity distributions in models of 10 day Cepheids
a) low mass (4.0 M₀) and b) high mass (7.4 M₀) using the
DYN code and KingIVA opacities.

Now it is clear that simply eyeballing the resulting light curves for comparison to observations is highly subjective (Simon and Davis 1983) but for these long period Cepheids the spiked peak makes Fourier analysis difficult. It is also possible that the so called "bumps" or shoulders on the rising branch of LP Cepheid light curves could be the "dips" now resolved by Moffett and Barnes. These "dips" appear to be due to a surface phenomena that is more easily described by the methods of dynamic zoning used in DYN. Even though the shock is treated by the use of "Pseudo-viscosity" it is well resolved by the fine zones included in the shock forming region. In the atmosphere of the star the strongest shock that develops is the one caused by the stopping of the infall of the envelope near the phase of maximum inward velocity. This shock gives rise to the "artificial viscosity dip" discussed by Davis (1975) and others. In most models another shock develops at the time of rapid expansion. This shock is associated with the brief transition of the ionization front from a "D" to a "R" type (Kahn's notation see Castor and Adams 1974) as it moves rapidly inward in mass. This inward shock remains below optical depth unity and therefore should not effect the light curves. Without the capability to follow line transport in a moving media we must rely on the photospheric continuum results for our comparisons to the observations.




Fig. 2 - The Dyn non-linear long period Cepheid model results with 60X evolutionary mass (left panel) and near evolutionary mass (right panel) compared to the observations of Moffett and Barnes (center panel).

THE RESULTS AND CEPHEID MASSES

In modeling the non-linear hydrodynamics that occurs in pulsating stars we have considered the effects of radiative transfer but not the effect of convection. It is believed that convection causes the limit to the red edge of the instability strip but is not very important to the light output otherwise. For long period Cepheids (observe Fig. 2) we generally find a "dip" in the rising part of the light curve which signals "shock dissipation" and possibly the limit on the amplitude of the pulsation. These "dips" may have previously been mistaken for "bumps" or shoulders and therefore the resonance effect. As discussed in our paper on X Cygni to remove the bumps on the light curves we need to use masses near the evolutionary masses. More and detailed velocity measurements, in synchronism with the light measurements, may help answer the question, "Are there "bumps" as well as "dips" in the long period Cepheid light curves?"

REFERENCES

- Adams, T.F. and Castor, J.I., Ap. J. 230, 826.
Castor, J.I., Davidson, D.K., and Davis, C.G., LA-6664.
Christy, R.F. (1968). Quart. J.R. A.S. 171, 593.
Connolly, L.P., (1980). Pub. A.S.P. 92, 165.
Cox, A.M., Brownlee, R.R. and Eilers, D.D. (1966). Ap. J. 144, 1024.
Cox, J.P. (1980) Theory of Stellar Pulsation, Princeton, Princeton University Press).
Davis, C.G. (1975) Cepheid Modeling, NASA SP-383.
Davis, C.G. Moffett, T.J. and Barnes, T.G. (1981) Ap. J. 246, 914.
Keller, C. and Mutschlecner, J.P. (1972) Ap. J. 171, 593.
Moffett, T.J. and Barnes, T.G., 1984 Ap. J. Suppl. in press.
Schmidt, E.G. and Parsons, S.B. (1982) Ap. J. 48, 165..
Simon, N.R. and Davis, C.G. (1983) Ap. J. 266, 787.
Simon, N.R. (1984) to be published.
Takeuti, M., UJI-IVE, K. and Aikawa, T. (1983) Sei. Rept. Tohoku University Vol. 14. No. 1.
Whitnev, C. (1983) Ap. J. 274, 830.